arXiv:1212.0097v1 [nlin.SI] 1 Dec 2012

Mixed solitons in $(2+1)$ dimensional multicomponent long wave-short wave system

T. Kanna*
Post Graduate and Research Department of Physics,
Bishop Heber College, Tiruchirapalli–620 017, India.

M. Vijayajayanthi
Department of Physics, Anna University, Chennai–600 025, India.

M. Lakshmanan†
Centre for Nonlinear Dynamics, School of Physics,
Bharathidasan University, Tiruchirapalli–620 024, India.

(Dated: December 4, 2012)

Abstract

We derive a (2+1)-dimensional multicomponent long wave-short wave resonance interaction (LSRI) system as the evolution equation for propagation of N-dispersive waves in weak Kerr type nonlinear medium in the small amplitude limit, by following the lines of reference [14] for the two-component case. The mixed (bright-dark) type soliton solutions of a particular (2+1)-dimensional multicomponent LSRI system, deduced from the general multicomponent higher dimensional LSRI system, are obtained by applying the Hirota's bilinearization method. Particularly, we show that the solitons in the LSRI system with two short-wave components behave like scalar solitons. We demonstrate that the bright solitons in the N-component LSRI system with $N > 2$ display interesting collision behaviour resulting in energy exchange among the bright solitons, whereas the dark solitons undergo standard elastic collision accompanied by a position shift and a phase-shift. Our analysis on the mixed bound solitons shows that the additional degree of freedom resulting due to the higher dimensional nature of the system results in a wide range of parameters for which the soliton collision can take place.

---

* Corresponding author Electronic mail: kanna_phy@bhc.edu.in.
† Electronic mail: lakshman@cnld.bdu.ac.in.

I. INTRODUCTION

The nonlinear interaction of multiple waves results in several new physical processes [1–3]. It has been shown in the two layer fluid model that resonance between the long-wave component and short-wave component occurs when the phase velocity of the former matches the group velocity of the latter [4]. This is a ubiquitous phenomenon which appears in hydrodynamics [5], bio-physics [6], plasma physics [7], and in nonlinear optical systems [8]. Though there are many studies on the long wave-short wave resonance interaction (LSRI) in one dimension [7–12], results are scarce for multicomponent higher dimensional LSRI system. In the context of nonlinear optics, the interaction of bright and small amplitude dark pulses in optical fiber is governed by the integrable Zakharov model [7, 8].

The resonance interaction of long-wave with short-wave was first investigated by Benney for capillary-gravity waves in deep water [13]. In this case simple interaction equations cannot be obtained, because for deep water waves there is no wave in the long wavelength limit. However, simple interaction equations can be deduced in a stable stratified fluid for oblique propagation of long and short-waves [5]. The single component two-dimensional LSRI equation for a two-layer fluid model has been derived in Ref. [4] by using the multiple scale perturbation method and bright and dark type one- and two-soliton solutions have been reported. In Ref. [14], Ohta *et al* derived the integrable two component analogue of the two-dimensional LSRI system as a governing equation for the interaction of the nonlinear dispersive waves by applying the reductive perturbation method. Very recently, the non-integrable three component Gross-Pitaevskii equations have been reduced to single component Yajima-Oikawa system by using multiple scale method [15]. In another recent work [16], the one-dimensional integrable two-component Zakharov-Yajima-Oikawa equation has been derived using multiple scale method and special bright-dark one-soliton solutions have been reported.

In Ref. [17], we have obtained general bright M-soliton solution, for arbitrary M, of the same integrable multicomponent $(2+1)$D-LSRI system (see Eq. (12)) considered in the present paper. Our earlier work [17] on bright soliton solutions of the multicomponent LSRI system shows that the role of long interfacial wave is to induce nonlinear interaction among the short-wave components resulting in non-trivial (shape-changing) collision behaviour characterized by energy exchange among the two short-wave components. As far as we know, for the first time in Ref. [17] we have identified the shape-changing/energy sharing collisions of bright solitons in a two dimensional integrable nonlinear system. This system will act as a potential candidate for realizing soliton collision-based computing and multistate logic [18–20]. Now it is of interest to derive the general two dimensional N-component equations describing the interaction of several short-wave packets with long-waves in a physical set up and to look for other types of multicomponent soliton solutions.

In recent years, much attention has been paid to investigate mixed (bright-dark) soliton dynamics in different dynamical systems including nonlinear optical systems and Bose-Einstein condensates [1, 8, 14–16, 21–23] of coupled bright-dark solitons and to analyse their propagation properties and collision dynamics. In the present work, we derive the (2+1)-dimensional N-component LSRI system governing the evolution of m short-waves and n long-waves with $(m+n=N)$ in a nonlinear dispersive medium and reduce the system to an integrable system for a particular choice of the system parameters. Then by applying the elegant Hirota's direct method to the integrable multicomponent LSRI system which results for particular choice, we obtain the coupled bright-dark one- and two- soliton solutions. We will show that bright and dark parts of the mixed solitons in the two short-wave components case behave like scalar solitons whereas if we go for three and more short-wave components the multicomponent nature of the solitons will come into picture and one can observe interesting propagation and collision properties. It is straight-forward albeit lengthy procedure to extend the analysis to construct M-soliton solution, with arbitrary M.

The present paper is organized as below. The general N-component LSRI system is derived by applying the multiple scale perturbation method in the next section. In section III, bilinear equations for the integrable (2+1)-dimensional multicomponent LSRI system are given. Sections IV and V deal with the mixed one- and two-soliton solutions of the multicomponent LSRI system. The collision dynamics of the solitons are discussed in section VI. Discussion on the mixed soliton bound states is presented in section VII. The final section is allotted for conclusion.

II. THE MODEL

To start with we obtain the general two-dimensional multicomponent evolution equation for the propagation of N-dispersive waves in a Kerr type nonlinear medium (ex.: optical fiber, photo-refractive medium) by generalizing the approach developed in refs. [8, 14] for the two and three components case. The waves are assumed to obey the following weakly nonlinear dispersion relations

$$\omega_j = \omega_j(\mathbb{K}_j; \mathbb{L}_j : |A_1|^2, |A_2|^2, ..., |A_N|^2), \quad j = 1, 2, 3, ..., N, \tag{1}$$

where $\mathbb{K}_j$ and $\mathbb{L}_j$ are the x and y components of the wave vector, $A_j(\equiv A_j(x, y, t))$ and ω_j are the complex amplitude and angular frequency of the j-th wave. The fundamental carrier wave is of the form $e^{i(\mathbb{K}_0 x+\mathbb{L}_0 y-\omega_0 t)}$. The most convenient way to derive the evolution equation for the amplitudes A_j's is to Taylor expand the angular frequencies ω_j's around the x and y components of the wave vector of the carrier wave $\mathbb{K}_0$ and $\mathbb{L}_0$, respectively, and the central frequency ω_0 at $|A_j| = 0$, as below:

$$\begin{aligned}(\omega_j - \omega_0) \;=\; & (\omega_{j,\mathbb{K}_j})_0 \Delta\mathbb{K}_j + (\omega_{j,\mathbb{L}_j})_0 \Delta\mathbb{L}_j + \frac{1}{2}(\omega_{j,\mathbb{K}_j\mathbb{K}_j})_0 \Delta\mathbb{K}_j^2 + \frac{1}{2}(\omega_{j,\mathbb{L}_j\mathbb{L}_j})_0 \Delta\mathbb{L}_j^2 \\ & +(\omega_{j,\mathbb{K}_j\mathbb{L}_j})_0 (\Delta\mathbb{K}_j)(\Delta\mathbb{L}_j) + \sum_{m=1}^{N} \left(\omega_{j,|A_m|^2}\right)_0 |A_m|^2 + ..., \quad j = 1, 2, ..., N,\end{aligned} \tag{2}$$

where $\Delta\mathbb{K}_j = \mathbb{K}_j - \mathbb{K}_0$, $\Delta\mathbb{L}_j = \mathbb{L}_j - \mathbb{L}_0$, $j = 1, 2, ..., N$. In this Taylor expansion and in the following, the subscript '0' in the $(\ )_0$ represents that the quantity appearing inside the bracket is evaluated at $\mathbb{K}_j = \mathbb{K}_0$, $\mathbb{L}_j = \mathbb{L}_0$, $\omega_j = \omega_0$ and $|A_j| = 0$. In Eq. (2), $\omega_{j,\mathbb{K}_j} = \frac{\partial\omega_j}{\partial\mathbb{K}_j}$, $\omega_{j,\mathbb{L}_j} = \frac{\partial\omega_j}{\partial\mathbb{L}_j}$, $\omega_{j,\mathbb{K}_j\mathbb{K}_j} = \frac{\partial^2\omega_j}{\partial\mathbb{K}_j^2}$, $\omega_{j,\mathbb{L}_j\mathbb{L}_j} = \frac{\partial^2\omega_j}{\partial\mathbb{L}_j^2}$, $\omega_{j,\mathbb{K}_j\mathbb{L}_j} = \frac{\partial^2\omega_j}{\partial\mathbb{K}_j\partial\mathbb{L}_j}$ and $\omega_{j,|A_m|^2} = \frac{\partial\omega_j}{\partial|A_m|^2}$. Then by replacing $(\omega_j - \omega_0)$, $\Delta\mathbb{K}_j$ and $\Delta\mathbb{L}_j$ by the operators $-i\frac{\partial}{\partial t}$, $i\frac{\partial}{\partial x}$ and $i\frac{\partial}{\partial y}$, respectively and transforming to the moving co-ordinates $x' = x - \overline{\omega}_1 t$, $y' = y - \overline{\omega}_2 t$, $t' = t$, with the assumption that beyond a particular component (say q^{th}) all the derivatives $\left(\frac{\partial\omega_j}{\partial\mathbb{K}_j}\right)_0$, $j = q+1, q+2, ..., N$, are same and so also the derivatives $\left(\frac{\partial\omega_j}{\partial\mathbb{L}_j}\right)_0$, i.e., $\left(\omega_{q+1,\mathbb{K}_{q+1}}\right)_0 \left(\equiv \frac{\partial\omega_{q+1}}{\partial\mathbb{K}_{q+1}}\right) = \left(\omega_{q+2,\mathbb{K}_{q+2}}\right)_0 = ... = \left(\omega_{N,\mathbb{K}_N}\right)_0 \equiv \overline{\omega_1}$ (say) and $\left(\omega_{q+1,\mathbb{L}_{q+1}}\right)_0 = \left(\omega_{q+2,\mathbb{L}_{q+2}}\right)_0 = ... = \left(\omega_{N,\mathbb{L}_N}\right)_0 \equiv \overline{\omega_2}$ (say), and omitting the primes for simplicity of notation we get

$$iA_{j,t} \;+\; iv_{jx}A_{j,x} + iv_{jy}A_{j,y} + C_1^{(j)}A_{j,xx} + C_2^{(j)}A_{j,yy} + C_3^{(j)}A_{j,xy} + \sum_{l=1}^{N} B_l^{(j)}|A_l|^2 A_j = 0, \tag{3a}$$

$$iA_{p,t} \;+\; C_4A_{p,xx} + C_5A_{p,yy} + C_6A_{p,xy} + \sum_{l=1}^{N} B_l^{(p)}|A_l|^2 A_p = 0, \tag{3b}$$

$$j = 1, 2, ..., q, \quad p = q+1, q+2, ..., N.$$

Here the independent variables appearing in the suffixes after the comma denote partial derivatives with respect to that variables and the group velocities of the j^{th} component along the x and y directions are $v_{jx} = ((\omega_{j,\mathbb{K}_j})_0 - \overline{\omega}_1)$ and $v_{jy} = ((\omega_{j,\mathbb{L}_j})_0 - \overline{\omega}_2)$, respectively. Various quantities in the above equations (3) are defined as $C_1^{(j)} = \left(-\frac{\omega_{j,\mathbb{K}_j\mathbb{K}_j}}{2}\right)_0$, $C_2^{(j)} = \left(-\frac{\omega_{j,\mathbb{L}_j\mathbb{L}_j}}{2}\right)_0$, $C_3^{(j)} = \left(-\omega_{j,\mathbb{K}_j\mathbb{L}_j}\right)_0$, $C_4 = \left(\frac{-\overline{\omega}_{1,\mathbb{K}_N}}{2}\right)_0$, $C_5 = \left(\frac{-\overline{\omega}_{2,\mathbb{L}_N}}{2}\right)_0$, $C_6 = (-\overline{\omega}_{1,\mathbb{L}_N})_0$, $B_j^{(i)} = \left(\frac{\partial \omega_i}{\partial |A_j|^2}\right)_0 \equiv \left(\omega_{i,|A_j|^2}\right)_0$, $i, j = 1, 2, 3, ..., N$.

We consider the case where the first q-components are in the anomalous dispersion region and the remaining $(N-q)$-components are in the normal dispersion regime. Following Ref. [8], the solutions of (3) are sought in the form

$$A_j = \psi_j(x,y,t)e^{i\delta_j t}, \quad j = 1, 2, ..., q, \tag{4a}$$

$$A_p = (u_0 + a_p(x,y,t))e^{i[\Lambda_p t + \vartheta_p(x,t)]}, \quad p = q+1, q+2, ..., N, \tag{4b}$$

where $\delta_j = \sum_{l=q+1}^{N} B_l^{(j)} u_0^2$, $\Lambda_p = \sum_{l=q+1}^{N} B_l^{(p)} u_0^2$ and a_p's are assumed to take only small values.

Substituting equations (4) in (3a) and neglecting the higher order terms involving (a_p, ϑ_p) and also their derivatives result in the equation

$$\begin{aligned} & i(\psi_{j,t} + v_{jx}\psi_{j,x} + v_{jy}\psi_{j,y}) + C_1^{(j)}\psi_{j,xx} + C_2^{(j)}\psi_{j,yy} + C_3^{(j)}\psi_{j,xy} \\ & \quad + \left(\sum_{l=1}^{q} B_l^{(j)}|\psi_l|^2\right)\psi_j + \left(\sum_{p=q+1}^{N} B_p^{(j)}(2u_0 a_p)\right)\psi_j = 0, \quad j = 1, 2, ..., q. \end{aligned} \tag{5}$$

In a similar manner, by incorporating (4) in (3b) and collecting the real and imaginary parts, we arrive at a set of two coupled equations. The resulting coupled equations can be grouped together to the following equation by differentiating the imaginary part equation twice with respect to 't' and making use of the real part equation:

$$\begin{aligned} & a_{p,tt} \ + \ C_4^2 a_{p,xxxx} + C_4 C_5 a_{p,xxyy} + C_4 C_6 a_{p,xxxy} + C_4 u_0 \left(\sum_{j=1}^{q} B_j^{(p)}|\psi_j|^2_{xx}\right) \\ & \quad + 2C_4 u_0^2 \sum_{l=q+1}^{N} B_l^{(p)} a_{l,xx} = 0, \quad p = q+1, q+2, ..., N. \end{aligned} \tag{6}$$

Equations (5) and (6) are general equations describing the two dimensional propagation of q waves in anomalous dispersion region and $(N-q)$-waves in the normal dispersion region. For $N = 3$ case with $q = 2$ the system has two-SW components and one long-wave component and coincides with the corresponding equations presented in Ref. [14]. One can notice from the general form of equations (5) and (6) that for the same $N(=3)$ but with different q value, (say $q = 1$), there is another possibility which will have one SW component and two long-wave components, and ultimately the dynamics will be different from the $q = 2$ case. Also, this systematic generalization to N component case is necessary to identify the way by which the additional wave components (modes) in the normal dispersion regime alter the governing equation for the three components case given in Ref. [14].

To deduce an integrable equation associated with the combined systems (5) and (6) we choose all the $B_l^{(p)}$'s, $l, p = q+1, ..., N$, in Eq. (6) to be equal to a constant value (say $-\gamma_1$, $\gamma_1 > 0$). Then we get

$$a_{p,tt} \;+\; C_4^2 a_{p,xxxx} + C_4C_5 a_{p,xxyy} + C_4C_6 a_{p,xxxy} + C_4 u_0 \left(\sum_{j=1}^{q} B_j^{(p)} |\psi_j|^2_{xx}\right)$$

$$-2C_4\gamma_1 u_0^2 \sum_{l=q+1}^{N} a_{l,xx} = 0, \quad p = q+1, q+2, ..., N. \tag{7}$$

In the following, we investigate the cumulative effect of the small amplitudes a_p's on the SW components by considering the superposition of these amplitudes involving only the sum of all the amplitudes and neglect all other combinations as they will be small due to the smallness of a_p's. Particularly, we add all the a-equations and define $\sum_{l=q+1}^{N} a_l = L$. By doing so we get

$$L_{tt} + C_4^2 L_{xxxx} + C_4C_5 L_{xxyy} + C_4C_6 L_{xxxy} + C_4 u_0 \left(\sum_{p=q+1}^{N}\sum_{j=1}^{q} B_j^{(p)} |\psi_j|^2_{xx}\right)$$

$$-2C_4[N-q]u_0^2\gamma_1 L_{xx} = 0. \tag{8}$$

The dispersion relation for the linear excitation corresponding to the long-wave components is found as

$$\Omega^2 = c^2k^2\left[1 + \frac{C_4^2}{c^2}k^2 + \frac{C_4C_5}{c^2}l^2 + \frac{C_4C_6}{c^2}kl\right], \tag{9}$$

where $c^2 = 2C_4u_0^2(N-q)\gamma_1$. Note that the velocity of the linear excitation depend upon the number of components N and increases as we increase the number of components (modes) in the normal dispersion regime. Thus our systematic generalization to N-component case shows that by altering the number of components in the normal dispersion region one can change the velocity of the pulse.

Next we apply the multiple scale approximation method to derive the two-dimensional multicomponent LSRI system as in Ref. [14]. We re-scale the variables t, x, y, a_p and ψ_j as

$$t'' = \epsilon t, \quad x'' = \sqrt{\epsilon}(x+ct), \quad y'' = \epsilon y, \quad a_p = \epsilon\,\hat{a}_p, \;\; \psi_j = \epsilon^{3/4}S^{(j)}, \tag{10}$$

where c is as defined above after the dispersion relation (9). Then the following set of equations results from Eq. (8) at the order $\epsilon^{5/2}$

$$2cL_{xt} + C_4 u_0 \left(\sum_{p=q+1}^{N} \sum_{j=1}^{q} B_j^{(p)} |S^{(j)}|^2_{xx} \right) = 0, \quad p = q+1, q+2, ..., N. \tag{11a}$$

At the order of $\epsilon^{5/4}$, we notice from Eq. (5) that all the group velocities of the short-wave components along the x direction are the same and their magnitudes are equal to the phase velocity of the long-wave component 'c' (i.e. $v_{jx} = -c, \ j = 1, 2, ..., q$). This is the condition for resonant interaction between long-waves and short-waves. Equation (5) reduces to the following set of coupled equations at the order of $\epsilon^{7/4}$ after replacing v_{jx} by c and rescaling of the variables as defined in equation(10),

$$i\left(S_t^{(j)} + v_{jy} S_y^{(j)}\right) + C_1^{(j)} S_{xx}^{(j)} + \left(2u_0 \sum_{p=q+1}^{N} B_p^{(j)} \hat{a}_p\right) S^{(j)} = 0, \quad j = 1, 2, ..., q. \tag{11b}$$

In Eq. (11), after applying the transformations (10) the double primes in the new variables 't', 'x' and 'y' are dropped, for convenience.

Equation (11) is the multicomponent LSRI system in $(2+1)$-dimensions which is non-integrable in general. By suitably choosing the constants $B_j^{(p)}$'s, $B_p^{(j)}$'s, C_4, $C_1^{(j)}$, $j = 1, 2, ..., q$, $p = q+1, ..., N$, and γ_1, along with the assumption that there is no group velocity delay between the short-wave components, we arrive at the following $(q+1)$-component (2+1)-dimensional LSRI system for the 2-dimensional propagation of N dispersive waves in weak Kerr-like nonlinear media,

$$i\left(S_t^{(j)} + S_y^{(j)}\right) - S_{xx}^{(j)} + LS^{(j)} = 0, \quad j = 1, 2, ..., q, \tag{12a}$$

$$L_t = 2\sum_{j=1}^{q} |S^{(j)}|^2_x. \tag{12b}$$

In Eq. (12), the subscripts denote the partial derivatives with respect to those independent variables. As mentioned in the introduction, we have obtained more general bright M-soliton solution, with arbitrary M, of Eq. (12) [17]. In reference [17], we have expressed the bright M-soliton solution of (12) in Gram determinant form and explicitly proved that the general multisoliton solution indeed satisfies the bilinear equations. We have also pointed out in the same work that for the two-SW components case ($q = 2$), the bright soliton solutions reported by Ohta *et al* in Ref. [14] follow as special cases of our general multi-soliton solutions [17]. As the $(q+1)$-component LSRI system (12) admits M-soliton solution, for arbitrary M [17], the system can be integrable [24, 25]. The study on the other integrable aspects of Eq. (12) is under progress and will be published elsewhere.

III. HIROTA'S BILINEARIZATION METHOD FOR THE $(2+1)D$ MULTICOMPONENT LSRI SYSTEM

Hirota's direct method is an elegant and powerful tool to construct the soliton solutions of a given integrable nonlinear evolution equation [24]. We look for the coupled bright-dark (mixed) soliton solutions of Eq. (12) in which the mixed soliton has m bright parts and n dark parts, such that $m+n=q$. To construct the mixed type soliton solutions, we perform the bilinearizing transformations, $S^{(j)}=\frac{g^{(j)}}{f}$, $S^{(m+l)}=\frac{h^{(l)}}{f}$, $L=-2\frac{\partial^2}{\partial x^2}(\ln f)$, $j=1,2,\ldots,m$, and $l=1,2,\ldots,n$, $(m+n=q)$, where $g^{(j)}$'s and $h^{(l)}$'s are arbitrary complex functions of x,y and t while f is a real function. The resulting bilinear equations are

$$D_1\left(g^{(j)}\cdot f\right)=0, \quad j=1,2,\ldots,m, \tag{13a}$$

$$D_1\left(h^{(l)}\cdot f\right)=0, \quad l=1,2,\ldots,n, \tag{13b}$$

$$D_2(f\cdot f)=-2\left(\sum_{j=1}^{m}g^{(j)}g^{(j)*}+\sum_{l=1}^{n}h^{(l)}h^{(l)*}\right), \tag{13c}$$

where $D_1=i(D_t+D_y)-D_x^2$ and $D_2=(D_tD_x-2\lambda)$, D-s are the standard Hirota's bilinear operators [24], $*$ stands for complex conjugation and λ is a constant yet to be determined. One can have bright solitons for the choice $\lambda=0$ in D_2 [14, 17] and the bright soliton collision dynamics of system (12) has been discussed in Ref. [17]. However for non-vanishing 'λ' values, the system can admit coupled bright-dark and dark-dark type soliton solutions. In this paper, we focus only on mixed (bright-dark) solitons corresponding to mixed type boundary conditions, that is, $S^{(j)},\ L \overset{x,y,t\to\pm\infty}{\longrightarrow} 0$, $S^{(m+l)} \overset{x,y,t\to\pm\infty}{\longrightarrow}$ constant, $j=1,2,...,m,\ l=1,2,...,n$.

This procedure can be very well applied to construct the dark-dark soliton solutions also. Here for convenience we consider the first 'm' short-wave components to be comprised of bright parts of the mixed solitons and the remaining $n(\equiv q-m)$ components to exhibit dark parts of the mixed solitons. To construct the mixed soliton solutions we expand the variables $g^{(j)}$'s, $h^{(l)}$'s and f as power series expansions in a standard way [23, 24]. After solving the resultant set of equations recursively we can obtain the explicit forms of $g^{(j)}$'s, $h^{(l)}$'s and f and hence the multisoliton solutions can be constructed.

IV. MULTICOMPONENT MIXED TYPE ONE-SOLITON SOLUTION

The mixed one-soliton solution of (12) with m-bright and n-dark parts can be obtained by restricting the power series expansions as $g^{(j)} = \chi g_1^{(j)}$, $h^{(l)} = h_0^{(l)}(1+\chi^2 h_2^{(l)})$, $f = 1+\chi^2 f_2$ and by solving the resulting equations, after their substitution into the bilinear equations (13) at various powers of χ recursively. The mixed one-soliton solution can be expressed in the following standard form,

$$S^{(j)} = A_j k_{1R} \,\text{sech}\left(\eta_{1R} + \frac{R}{2}\right) e^{i\eta_{1I}}, \quad j = 1,2,...,m, \tag{14a}$$

$$S^{(m+l)} = \rho_l \, e^{i(\zeta_l+\phi_l+\pi)} \left[\cos(\phi_l) \tanh\left(\eta_{1R} + \frac{R}{2}\right) + i \,\sin(\phi_l)\right], \quad l = 1,2,...,n, \tag{14b}$$

$$L = -2k_{1R}^2 \,\text{sech}^2\left(\eta_{1R} + \frac{R}{2}\right). \tag{14c}$$

The various quantities appearing in the above equations are given below:

$$e^R = \frac{1}{4}\left(\sum_{j=1}^{m} |\alpha_1^{(j)}|^2\right)\left(\sum_{l=1}^{n} |\rho_l|^2 \cos^2(\phi_l) - \omega_{1R}k_{1R}\right)^{-1}, \quad A_j = \frac{\alpha_1^{(j)} e^{-\frac{R}{2}}}{(k_1+k_1^*)}, \tag{14d}$$

$$\phi_l = \tan^{-1}\left(\frac{k_{1I}-m_l}{k_{1R}}\right), \quad \eta_{1R} = k_{1R}\left[x + \left(2k_{1I} - \frac{\omega_{1R}}{k_{1R}}\right)y + \left(\frac{\omega_{1R}}{k_{1R}}\right)t\right], \tag{14e}$$

$$\eta_{1I} = k_{1I}x - (k_{1R}^2 - k_{1I}^2 + \omega_{1I})y + \omega_{1I}t, \quad \zeta_l = (m_l^2 - b_l)t + b_l y + m_l x. \tag{14f}$$

In equations (14) and in the following the suffixes R and I of a particular quantity denote the real and imaginary parts of that quantity, respectively. Also $\alpha_1^{(j)}$'s, $j = 1,2,...,m$, ρ_l, $\omega_1(=\omega_{1R}+i\omega_{1I})$ and $k_1 = k_{1R}+ik_{1I}$ are complex parameters, while m_l and b_l, $l = 1,2,...,n$, are real parameters. The above solution is non-singular for the condition $\sum_{l=1}^{n}|\rho_l|^2\cos^2(\phi_l) > \omega_{1R}k_{1R}$. The amplitude (peak value) of the j^{th} bright part of the mixed soliton is $A_j k_{1R}$ and that of $(m+l)$-th dark part of the mixed soliton is ρ_l. The speed of the soliton is ω_{1R}/k_{1R}. It can be noticed that both parts of the soliton have the same central position $R/2k_{1R}$. But their phases are different. In fact, the phase of the dark component has two contributions, one from the background carrier wave and the other from ϕ_l. The quantity ϕ_l indeed determines the darkness of the dark soliton. It is interesting to notice that the bright and dark parts of the mixed soliton of the LSRI system with more than two short-wave components display several interesting features in contrast to the case of just two-SW components as will be shown. To elucidate the understanding of such behaviour we present the explicit forms of one- soliton solutions for the two- and three- SW components and analyse them in the following subsections. For brevity, in the following we refer to mixed M-soliton solution with m-bright parts and n-dark parts as (mb-nd) mixed M soliton solution.

A. Two-SW components ($m = 1$, $n = 1$) case

This case admits only a simple type of bright-dark pair in which the bright part of mixed soliton appears in the first component and the dark part of the mixed soliton in the remaining component or vice-versa. The one-soliton solution for this case can be expressed as

$$S^{(1)} = \left(\sqrt{|\rho_1|^2 \cos^2(\phi_1) - \omega_{1R} k_{1R}}\right) \operatorname{sech}\left(\eta_{1R} + \frac{R}{2}\right) e^{i(\eta_{1I}+\theta)}, \tag{15a}$$

$$S^{(2)} = \rho_1 \, e^{i(\zeta_1+\phi_1+\pi)} \left[\cos(\phi_1) \tanh\left(\eta_{1R} + \frac{R}{2}\right) + i \sin(\phi_1)\right], \tag{15b}$$

$$L = -2k_{1R}^2 \operatorname{sech}^2\left(\eta_{1R} + \frac{R}{2}\right), \tag{15c}$$

where $\frac{R}{2} = \ln\left[\frac{|\alpha_1^{(1)}|}{2\sqrt{|\rho_1|^2 \cos^2(\phi_1) - \omega_{1R} k_{1R}}}\right]$, $\phi_1 = \tan^{-1}\left(\frac{k_{1I}-m_1}{k_{1R}}\right)$, $\theta = \tan^{-1}\left(\frac{\alpha_{1I}^{(1)}}{\alpha_{1R}^{(1)}}\right)$, $\zeta_1 = (m_1^2 - b_1)t + b_1 y + m_1 x$, and η_{1R} and η_{1I} are given in equations (14e) and (14f).

The amplitude of the bright part $\left(\sqrt{|\rho_1|^2 \cos^2(\phi_1) - \omega_{1R} k_{1R}}\right)$ is independent of the parameter $\alpha_1^{(1)}$, but it is influenced significantly by the background carrier wave (ρ_1). Such a mixed soliton at $t = -3$ and $y = -1$ is depicted in Fig. 1 for the parametric choice $k_1 = -3 + i$, $\omega_1 = 1 + 0.7i$, $m_1 = 1.4$, $\rho = 1 - i$, $b_1 = -0.2$, $\alpha_1^{(1)} = 1 + i$.

B. Three-SW components ($q = 3$) case

Next we consider Eq. (12) with $q = 3$. For this case the mixed soliton can be split up into bright and dark parts among the three-SW components in two different ways. One corresponds to the (2b-1d) mixed soliton case where bright parts are in the $S^{(1)}$ and $S^{(2)}$ components while the dark part of mixed soliton appears in the $S^{(3)}$ component. The other possibility is a (1b-2d) mixed soliton case in which the bright part of the mixed



FIG. 1: Mixed one-soliton in two-component LSRI system. ($|S^{(1)}|^2$ - solid curve, $|S^{(2)}|^2$ - dashed curve)

soliton appears in the $S^{(1)}$ component while the dark parts are split among the remaining components $S^{(2)}$ and $S^{(3)}$.

1. *(2b-1d) mixed one-soliton solution*

The one-soliton solution for this case where the bright parts appear in the $S^{(1)}$ and $S^{(2)}$ components while the third component $S^{(3)}$ comprises the dark part of the mixed soliton, can be written from (14) as

$$S^{(j)} = A_j k_{1R} \,\text{sech}\left(\eta_{1R} + \frac{R}{2}\right) e^{i\eta_{1I}}, \quad j = 1, 2, \tag{16a}$$

$$S^{(3)} = \rho_1 \, e^{i(\zeta_1+\phi_1+\pi)} \left[\cos(\phi_1) \tanh\left(\eta_{1R} + \frac{R}{2}\right) + i \sin(\phi_1)\right], \tag{16b}$$

where $A_j = \left(\frac{\alpha_1^{(j)}}{2k_{1R}}\right) e^{-\frac{R}{2}}$, $j = 1, 2$, $\zeta_1 = (m_1^2 - b_1)t + b_1 y + m_1 x$, $R = \ln\left[\frac{|\alpha_1^{(1)}|^2 + |\alpha_1^{(2)}|^2}{4\left(|\rho_1|^2 \cos^2(\phi_1) - \omega_{1R} k_{1R}\right)}\right]$, $\phi_1 = \tan^{-1}\left(\frac{k_{1I} - m_1}{k_{1R}}\right)$, and η_{1R} and η_{1I} are as defined in eqn. (14). L takes the same form as in eqn. (15c) with the above redefinition of R. Here one can observe that the $\alpha_1^{(j)}$-parameters appear explicitly in the amplitude of the bright soliton. The (2b-1d) one-soliton solution is characterized by twelve real parameters $\alpha_{1R}^{(1)}$, $\alpha_{1I}^{(1)}$, $\alpha_{1R}^{(2)}$, $\alpha_{1I}^{(2)}$, k_{1R}, k_{1I}, ω_{1R}, ω_{1I}, ρ_{1R}, ρ_{1I}, m_1, and b_1 and is restricted by the condition $|\rho_1|^2 \cos^2(\phi_1) > \omega_{1R} k_{1R}$ for non-singular solutions. Such type of (2b-1d) bright one-soliton is shown in Fig. 2(a) at $t = -3$ and $y = -1$ for the parameters $k_1 = 2 - 2i$, $\omega_1 = -1 - i$, $m_1 = 3$, $\rho_1 = 1 - i$, $b_1 = -0.2$, $\alpha_1^{(1)} = 1.8 + i$, and $\alpha_1^{(2)} = 1$. One can also tune the intensity of bright parts without altering the depth of the dark part of the mixed soliton by suitably choosing the $\alpha_1^{(1)}$ parameter as can be evidenced from Fig. 2(b) which is drawn for same parameter as that of Fig. 2(a) except for $\alpha_1^{(1)} = 1+i$. The soliton appearing in the long-wave component looks similar in both the cases and so we do not present it here.



FIG. 2: Altering the intensity of bright soliton without affecting the dark soliton of (2b-1d) mixed one-soliton in three-component LSRI system by tuning $\alpha_1^{(j)}$ parameter. ($|S^{(1)}|^2$ - solid curve, $|S^{(2)}|^2$ - dashed curve, $|S^{(3)}|^2$ - dot-dashed curve)

2. *(1b-2d) mixed one-soliton solution*

This case corresponds to the appearance of the bright part of the mixed soliton in the $S^{(1)}$ component while its dark part appears in the $S^{(2)}$ and $S^{(3)}$ components. The corresponding mixed one-soliton solution is

$$S^{(1)} = \sqrt{|\rho_1|^2\cos^2(\phi_1) + |\rho_2|^2\cos^2(\phi_2) - \omega_{1R}k_{1R}}\ \operatorname{sech}\left(\eta_{1R} + \frac{R}{2}\right) e^{i(\eta_{1I}+\theta_1)}, \tag{17a}$$

$$S^{(1+l)} = \rho_l\, e^{i(\zeta_l+\phi_l+\pi)} \left[\cos(\phi_l)\tanh\left(\eta_{1R} + \frac{R}{2}\right) + i\sin(\phi_l)\right], \quad l = 1,2, \tag{17b}$$

$$L = -2k_{1R}^2 \operatorname{sech}^2\left(\eta_{1R} + \frac{R}{2}\right), \tag{17c}$$

where $R = \ln\left(\frac{|\alpha_1^{(1)}|^2}{(k_1+k_1^*)^2}\right) - \ln\left(|\rho_1|^2\cos^2(\phi_1) + |\rho_2|^2\cos^2(\phi_2) - \omega_{1R}k_{1R}\right)$, $\zeta_l = (m_l^2 - b_l)t + b_l y + m_l x$, $\phi_l = \tan^{-1}\left(\frac{k_{1I}-m_l}{k_{1R}}\right)$, $l = 1,2$, $\theta_1 = \tan^{-1}\left(\frac{\alpha_{1I}}{\alpha_{1R}}\right)$, and η_{1R} and η_{1I} are as in Eq. (14). This solution is characterized by five complex parameters $\alpha_1^{(1)}$, k_1, ρ_1, ρ_2 and ω_1 and four real parameters b_l and m_l, $l = 1,2$, along with the condition $|\rho_1|^2\cos^2(\phi_1) + |\rho_2|^2\cos^2(\phi_2) > \omega_{1R}k_{1R}$. It can be observed from the above solution that in contrast to the (2b-1d) case, here the amplitudes of the bright and dark parts cannot be controlled by the α parameters. For illustrative purpose, in Fig. 3 we have shown the (1b-2d) mixed one-soliton solution for the parameters $k_1 = 1 - i$, $\omega_1 = -1 - i$, $m_1 = 2$, $m_2 = -2$, $\rho_1 = 1 - i$, $\rho_2 = 1 + i$, $b_1 = -0.2$, $b_2 = 0.2$, and $\alpha_1^{(1)} = 0.2 - 0.01i$ at $t = -3$ and $y = -1$.



FIG. 3: (1b-2d) mixed one-soliton in three-component LSRI system.

V. MULTICOMPONENT MIXED TYPE TWO-SOLITON SOLUTIONS

It is a straightforward but lengthy procedure to construct the two-soliton solution. We obtain the mixed two-soliton solution of system (12) by restricting the power series expansion for $g^{(j)}$'s, $h^{(l)}$'s and f as $g^{(j)} = \chi g_1^{(j)} + \chi^3 g_3^{(j)}$, $h^{(l)} = h_0^{(l)}(1 + \chi^2 h_2^{(l)} + \chi^4 h_4^{(l)})$, $f = 1 + \chi^2 f_2 + \chi^4 f_4$, $j = 1, 2, \ldots, m,\ l = 1, 2, \ldots, n$, and following the standard procedure [23]. The explicit form of (mb-nd) mixed two-soliton solution is given below.

$$S^{(j)} = \frac{1}{D}\left(\alpha_1^{(j)} e^{\eta_1} + \alpha_2^{(j)} e^{\eta_2} + e^{\eta_1+\eta_1^*+\eta_2+\delta_{1j}} + e^{\eta_2+\eta_2^*+\eta_1+\delta_{2j}}\right), \qquad j = 1, 2, \ldots, m, \tag{18a}$$

$$\begin{aligned} S^{(l+m)} = \frac{1}{D}\Big[\rho_l\, e^{i\zeta_l}\Big(1 + e^{\eta_1+\eta_1^*+Q_{11}^{(l)}} + e^{\eta_1+\eta_2^*+Q_{12}^{(l)}} + e^{\eta_2+\eta_1^*+Q_{21}^{(l)}} \\ + e^{\eta_2+\eta_2^*+Q_{22}^{(l)}} + e^{\eta_1+\eta_1^*+\eta_2+\eta_2^*+Q_3^{(l)}}\Big)\Big], \qquad l = 1, 2, \ldots, n, \end{aligned} \tag{18b}$$

$$L = -2\frac{\partial^2}{\partial x^2}\left(\ln(D)\right), \tag{18c}$$

where

$$D = 1+e^{\eta_1+\eta_1^*+R_1}+e^{\eta_1+\eta_2^*+\delta_0}+e^{\eta_2+\eta_1^*+\delta_0^*}+e^{\eta_2+\eta_2^*+R_2}+e^{\eta_1+\eta_1^*+\eta_2+\eta_2^*+R_3}, \tag{18d}$$

$$\eta_j = k_jx-(ik_j^2+\omega_j)y+\omega_jt, \quad j=1,2, \tag{18e}$$

$$e^{\delta_{1j}} = \frac{(k_1-k_2)(\alpha_1^{(j)}\kappa_{21}-\alpha_2^{(j)}\kappa_{11})}{(k_1+k_1^*)(k_2+k_1^*)}, \quad e^{\delta_{2j}} = \frac{(k_2-k_1)(\alpha_2^{(j)}\kappa_{12}-\alpha_1^{(j)}\kappa_{22})}{(k_2+k_2^*)(k_1+k_2^*)}, \tag{18f}$$

$$e^{Q_{ij}^{(l)}} = -\frac{(k_i-ib_l)}{(k_j^*+ib_l)}\mu_{ij}, \quad i,j=1,2, \quad e^{Q_3^{(l)}} = \left[\frac{(k_1-ib_l)(k_2-ib_l)}{(k_1^*+ib_l)(k_2^*+ib_l)}\right]e^{R_3}, \tag{18g}$$

$$e^{R_1} = \mu_{11},\ e^{R_2}=\mu_{22},\ e^{\delta_0}=\mu_{12},\ e^{\delta_0^*}=\mu_{21},\ e^{R_3}=\frac{|k_1-k_2|^2(\kappa_{11}\kappa_{22}-\kappa_{12}\kappa_{21})}{(k_1+k_1^*)|k_1+k_2^*|^2(k_2+k_2^*)}, \tag{18h}$$

$$\kappa_{ip} = \sum_{j=1}^{m}(\alpha_i^{(j)}\alpha_p^{(j)*})\left(\sum_{l=1}^{n}\frac{|\rho_l|^2(k_i+k_p^*)}{(k_i-im_l)(k_p^*+im_l)}-(\omega_i+\omega_p^*)\right)^{-1}, \quad \mu_{ip}=\frac{\kappa_{ip}}{(k_i+k_p^*)}, \tag{18i}$$

$$\zeta_l = (m_l^2-b_l)t+b_ly+m_lx, \quad i,p=1,2; \ \ j=1,2,...,m; \ \ l=1,2,...,n. \tag{18j}$$

We discuss below the two- and three- SW components cases to bring out certain interesting features of the multicomponent LSRI system with $q>2$.

A. Two-SW components case ($m=1, n=1$)

For this case, the mixed two-soliton solution with its bright part appearing in the $S^{(1)}$ component and the dark part appearing in the $S^{(2)}$ component can be obtained by putting $m=1$ and $n=1$ in Eq. (18). The resulting (1b-1d) mixed two-soliton solution is characterized by seven complex parameters ($\alpha_1^{(1)}, \alpha_2^{(1)}, k_1, k_2, \omega_1, \omega_2$ and ρ_1) and two real parameters (b_1 and m_1). This mixed two-soliton solution is restricted by the conditions $|\rho_1|^2(k_i+k_p^*) > (\omega_i+\omega_p^*)(k_i-im_1)(k_p^*+m_1)$, $i,p=1,2$, as in the case of the one-soliton solution.

B. Three-SW components case ($m=2, n=1$)

1. $(2b-1d)$ *mixed two-soliton solution*

First we write down the mixed two-soliton solution with its bright parts in the first two SW components $S^{(1)}$ and $S^{(2)}$ while the dark part appear in the third component $S^{(3)}$. For brevity, we mention the straight-forward procedure to write down the soliton solution from Eq. (18) instead of presenting the explicit cumbersome expressions. The explicit forms of bright part of the mixed solitons appearing in $S^{(1)}$ and $S^{(2)}$ can be obtained by putting $j = 1$ and $j = 2$, respectively, in Eq. (18a). The dark part of the mixed soliton solution in the $S^{(3)}$ component results from Eq. (18b) for $l = 1$. In a similar manner, the various quantities appearing in the (2b-1d) mixed two-soliton solution can also be obtained by choosing $m = 2$ and $n = 1$ in the expressions for μ_{ip} and κ_{ip}. The two-soliton solution in this case is characterized by twenty real parameters with the conditions $|\rho_1|^2(k_i + k_p^*) > (\omega_i + \omega_p^*)(k_i - im_1)(k_p^* + im_1)$, $i, p = 1, 2$.

2. $(1b - 2d)$ *mixed two-soliton solution*

Another possible split up for the three SW components case is to have the bright part of the mixed two-solitons in one component (say $S^{(1)}$) and the other two components ($S^{(2)}$ and $S^{(3)}$) comprise of dark parts. The obtained (1b-2d) mixed two-soliton solution can be deduced from Eq. (18) by putting $m = 1$ and $n = 2$. This mixed type two-soliton solution is characterized by sixteen real parameters and is restricted by the conditions $\sum_{l=1}^{2} \frac{|\rho_l|^2(k_i + k_l^*)}{(k_i - im_l)(k_l^* + im_l)} > (\omega_i + \omega_p^*)$, $i, p = 1, 2$, for obtaining non-singular solutions.

We wish to remark that our above analysis can be extended in a straight-forward way to construct three- as well as multi- soliton solutions. We have indeed obtained the mixed three-soliton solution, but we desist from presenting the solution here due to its cumbersome expression. Also, from the three-soliton solution we identify that the soliton collision is pair-wise and there is no multi-particle effects. Hence a detailed analysis of two-soliton collision is necessary as the higher order soliton interactions can be analyzed in terms of two-soliton collision.

VI. SOLITON INTERACTION

The multicomponent mixed type two-soliton solution presented in the preceding section contains all the information regarding the dynamics of two solitons in the $(2+1)D$ multicomponent LSRI system. To elucidate the understanding of the collision of mixed solitons, we present the detailed asymptotic analysis of the two and three component cases separately in this section. In particular, we study the interaction of solitons in the $x - y$ plane. A similar approach can also be very well applied to study the collision dynamics in the $x-t$ plane.

We choose the soliton parameters as $k_{1R}>0,\ k_{2R}>0,\ k_{1I}>k_{2I},\ \frac{k_{2R}}{k_{1R}}>\left|\frac{\omega_{2R}}{\omega_{1R}}\right|,\ \frac{k_{2R}k_{2I}}{k_{1R}k_{1I}}>\left|\frac{\omega_{2R}}{\omega_{1R}}\right|$, without loss of generality. For this choice, we find that for a fixed "t" if the soliton, say s_1, is localized along the straight line $\eta_{1R}=k_{1R}x+(2k_{1R}k_{1I}-\omega_{1R})y+\omega_{1R}t\simeq 0$, then η_{2R} will tend to $\pm\infty$ as $(x,y)\to\pm\infty$. Similarly, if the soliton, say s_2, is localized along the straight line $\eta_{2R}=k_{2R}x+(2k_{2R}k_{2I}-\omega_{2R})y+\omega_{2R}t\simeq 0$, then $\eta_{1R}\to\pm\infty$.

A. Two-SW components case

A careful analysis of the asymptotic forms of the mixed two-soliton solution for the two SW components case shows that the intensities of the bright and dark parts of the solitons before and after collision remain unaltered. There occurs only a position-shift in the bright and dark parts of the two colliding solitons. The asymptotic forms of the solitons in these regions are given below.

(i) Before Collision $(x,y\to-\infty)$:

Soliton s_1

$$S_1^{(1)-} \simeq A_1^{1-}\text{sech}\left(\eta_{1R}+\frac{R_1}{2}\right)e^{i\eta_{1I}}, \tag{19a}$$

$$S_1^{(2)-} \simeq \rho_1 e^{i(\zeta_1+\phi_1^{(1)}+\pi)}\left[\cos(\phi_1^{(1)})\tanh\left(\eta_{1R}+\frac{R_1}{2}\right)+i\,\sin(\phi_1^{(1)})\right], \tag{19b}$$

$$L \simeq -2k_{1R}\,\text{sech}^2\left(\eta_{1R}+\frac{R_1}{2}\right). \tag{19c}$$

Soliton s_2

$$S_2^{(1)-} \simeq A_1^{2-}\text{sech}\left(\eta_{2R}+\frac{R_3-R_1}{2}\right)e^{i\eta_{2I}}, \tag{19d}$$

$$S_2^{(2)-} \simeq \rho_1 e^{i(\zeta_1+\phi_1^{(2)}+2\phi_1^{(1)})}\left[\cos(\phi_1^{(2)})\,\tanh\left(\eta_{2R}+\frac{R_3-R_1}{2}\right)+i\,\sin(\phi_1^{(2)})\right], \tag{19e}$$

$$L \simeq -2k_{2R}\,\text{sech}^2\left(\eta_{2R}+\frac{R_3-R_1}{2}\right). \tag{19f}$$

(ii) After Collision ($x, y \to +\infty$):

Soliton s_1

$$
\begin{aligned}
S_1^{(1)+} &\simeq A_1^{1+} \mathrm{sech}\left(\eta_{1R} + \frac{R_3 - R_2}{2}\right) e^{i\eta_{1I}}, && \text{(19g)}\\
S_1^{(2)+} &\simeq \rho_1 e^{i(\zeta_1 + \phi_1^{(1)} + 2\phi_1^{(2)})}\left[\cos(\phi_1^{(1)})\,\tanh\left(\eta_{1R} + \frac{R_3 - R_2}{2}\right) + i\,\sin(\phi_1^{(1)})\right], && \text{(19h)}\\
L &\simeq -2k_{1R}\,\mathrm{sech}^2\left(\eta_{1R} + \frac{R_3 - R_2}{2}\right). && \text{(19i)}
\end{aligned}
$$

Soliton s_2

$$
\begin{aligned}
S_2^{(1)+} &\simeq A_1^{2+} \mathrm{sech}\left(\eta_{2R} + \frac{R_2}{2}\right) e^{i\eta_{2I}}, && \text{(19j)}\\
S_2^{(2)+} &\simeq \rho_1 e^{i(\zeta_1 + \phi_1^{(2)} + \pi)}\left[\cos(\phi_1^{(2)})\,\tanh\left(\eta_{2R} + \frac{R_2}{2}\right) + i\,\sin(\phi_1^{(2)})\right], && \text{(19k)}\\
L &\simeq -2k_{2R}\,\mathrm{sech}^2\left(\eta_{2R} + \frac{R_2}{2}\right), && \text{(19l)}
\end{aligned}
$$

where $\eta_j, \alpha_j^{(1)}$, $j = 1,2$, $\alpha_2^{(1)}$, R_1, R_2, R_3, δ_{11}, and δ_{21} are defined in Eq. (18), $\phi_1^{(l)} = \tan^{-1}(\frac{k_{lI} - m_1}{k_{lR}})$, $l = 1,2$.

The amplitudes of the bright parts of the solitons s_1 and s_2 before interaction (A_1^{1-}, A_1^{2-}) and their amplitudes after interaction (A_1^{1+}, A_1^{2+}) are given by $A_1^{1-} = \frac{\alpha_1^{(1)} e^{-\frac{R_1}{2}}}{2}$, $A_1^{2-} = \frac{e^{\delta_{11} - \frac{R_1 + R_3}{2}}}{2}$, $A_1^{1+} = \frac{e^{\delta_{21} - \frac{R_2 + R_3}{2}}}{2}$ and $A_1^{2+} = \frac{\alpha_2^{(1)} e^{-\frac{R_2}{2}}}{2}$. Here and in the following, the superscript (subscript) of A's denotes the soliton (component) number and $-$ ($+$) represents the soliton before (after) collision. Upon substitution of the corresponding expressions for R's and δ's we find that the intensities of the bright parts of the solitons s_1 and s_2 are same before and after collision. Similarly, we find the amplitudes of dark part of mixed solitons appearing in the $S^{(2)}$ component before and after interaction are same and are equal to ρ_1. The two colliding solitons s_1 and s_2 appearing in the bright component ($S^{(1)}$) also experience a position-shift of opposite sense whose magnitude is given by $|\Phi| = \left|\frac{R_3 - R_2 - R_1}{2}\right|$. Additionally, the dark soliton s_1 (s_2) experiences a phase-shift $2\phi_1^{(2)} - \pi$ ($-2\phi_1^{(1)} + \pi$).

Thus the mixed solitons in the SW components undergo elastic collision. The solitons in the long-wave component also undergo elastic collision with mere position-shift of magnitude $|\Phi|$. The elastic collision of solitons in the two component LSRI system is shown in Fig. 4 for the parametric choice $k_1 = 1 - i$, $k_2 = 1.5 + 0.75i$, $\omega_1 = -1 - 2i$, $\omega_2 = -0.75 + i$, $m_1 = 2$, $\rho_1 = 2$, $b_1 = 7$, $\alpha_1^{(1)} = 0.3 + i$, and $\alpha_2^{(1)} = 0.05 - i$ at $t = -1$.



FIG. 4: Elastic collision of (1b-1d) mixed solitons in two-component LSRI system.

B. Three-SW components case

This case can admit two types of soliton collisions as mentioned in section V B. The collision scenario depends upon the splitting of mixed solitons into bright and dark parts among the components and displays interesting dynamical behaviour. To illustrate this, we discuss the soliton collision for this (2b-1d) case in detail.

1. (2b-1d) soliton collisions

The asymptotic forms of solitons (s_1 and s_2) before and after collision can be deduced from the exact two-soliton solutions (18) by putting $m = 2$ and $n = 1$.

(i) Before Collision $(x, y \to -\infty)$:

Soliton s_1

$$\begin{aligned} S_1^{(j)-} &\simeq A_j^{1-}\text{sech}\left(\eta_{1R}+\frac{R_1}{2}\right)e^{i\eta_{1I}}, \quad j=1,2, && \text{(20a)}\\ S_1^{(3)-} &\simeq \rho_1 e^{i(\zeta_1+\phi_1^{(1)}+\pi)}\left[\cos(\phi_1^{(1)})\,\tanh\left(\eta_{1R}+\frac{R_1}{2}\right)+i\,\sin(\phi_1^{(1)})\right], && \text{(20b)}\\ L &\simeq -2k_{1R}\,\text{sech}^2\left(\eta_{1R}+\frac{R_1}{2}\right). && \text{(20c)} \end{aligned}$$

Soliton s_2

$$\begin{aligned} S_2^{(j)-} &\simeq A_j^{2-}\text{sech}\left(\eta_{2R}+\frac{R_3-R_1}{2}\right)e^{i\eta_{2I}}, \quad j=1,2, && \text{(20d)}\\ S_2^{(3)-} &\simeq \rho_1 e^{i(\zeta_1+\phi_1^{(2)}+2\phi_1^{(1)})}\left[\cos(\phi_1^{(2)})\,\tanh\left(\eta_{2R}+\frac{R_3-R_1}{2}\right)+i\,\sin(\phi_1^{(2)})\right], && \text{(20e)}\\ L &\simeq -2k_{2R}\,\text{sech}^2\left(\eta_{2R}+\frac{R_3-R_1}{2}\right). && \text{(20f)} \end{aligned}$$

(ii) After Collision ($x, y \to +\infty$):

Soliton s_1

$$\begin{aligned}
S_1^{(j)+} &\simeq A_j^{1+} \mathrm{sech}\left(\eta_{1R} + \frac{R_3 - R_2}{2}\right) e^{i\eta_{1I}}, \quad j = 1, 2, && \text{(20g)}\\
S_1^{(3)+} &\simeq \rho_1 e^{i(\zeta_1 + \phi_1^{(1)} + 2\phi_1^{(2)})} \left[\cos(\phi_1^{(1)}) \tanh\left(\eta_{1R} + \frac{R_3 - R_2}{2}\right) + i \sin(\phi_1^{(1)})\right], && \text{(20h)}\\
L &\simeq -2k_{1R} \,\mathrm{sech}^2\left(\eta_{1R} + \frac{R_3 - R_2}{2}\right). && \text{(20i)}
\end{aligned}$$

Soliton s_2

$$\begin{aligned}
S_2^{(j)+} &\simeq A_j^{2+} \mathrm{sech}\left(\eta_{2R} + \frac{R_2}{2}\right) e^{i\eta_{2I}}, \quad j = 1, 2, && \text{(20j)}\\
S_2^{(3)+} &\simeq \rho_1 e^{i(\zeta_1 + \phi_1^{(2)} + \pi)} \left[\cos(\phi_1^{(2)}) \tanh\left(\eta_{2R} + \frac{R_2}{2}\right) + i \sin(\phi_1^{(2)})\right], && \text{(20k)}\\
L &\simeq -2k_{2R} \,\mathrm{sech}^2\left(\eta_{2R} + \frac{R_2}{2}\right). && \text{(20l)}
\end{aligned}$$

The various quantities appearing in the above equations can be obtained from Eq. (18) by putting $m = 2$ and $n = 1$. Here, $\phi_1^{(l)} = \tan^{-1}(\frac{k_{lI} - m_l}{k_{lR}})$, $l = 1, 2$. We find that the amplitudes of the bright parts of the two solitons s_1 and s_2 before and after interaction are related through the relation,

$$A_j^{l+} = T_l^j A_j^{l-}, \qquad j = 1, 2, \quad l = 1, 2, \tag{21a}$$

where $A_j^{1-} = \frac{\alpha_1^{(j)}}{2} e^{-R_1/2}$, $A_j^{2-} = \frac{1}{2} e^{\delta_{1j} - \frac{R_1 + R_3}{2}}$ and the transition amplitudes T_l^j's are defined as

$$T_1^j = \left(\frac{k_2 - k_1}{k_2^* - k_1^*}\right)\left(\frac{k_1^* + k_2}{k_1 + k_2^*}\right)^{1/2} \left(\frac{\left(\frac{\alpha_2^{(j)}}{\alpha_1^{(j)}}\right)\lambda_1 - 1}{\sqrt{1 - \lambda_1\lambda_2}}\right), \tag{21b}$$

$$T_2^j = \left(\frac{k_1 - k_2}{k_1^* - k_2^*}\right)\left(\frac{k_1^* + k_2}{k_1 + k_2^*}\right)^{1/2} \left(\frac{\sqrt{1 - \lambda_1\lambda_2}}{\left(\frac{\alpha_1^{(j)}}{\alpha_2^{(j)}}\right)\lambda_2 - 1}\right), \; j = 1, 2, \tag{21c}$$

in which $\lambda_1 = \frac{\kappa_{12}}{\kappa_{22}}$ and $\lambda_2 = \frac{\kappa_{21}}{\kappa_{11}}$. Here the κ_{ip}'s can be obtained from Eq. (18) by putting $m = 2$ and $n = 1$. This shows that the intensities of the bright parts of the mixed solitons before and after collision are different in general. The transition amplitudes T_l^j's, $l, j = 1, 2$, become unimodular only for the choice $|\alpha_1^{(1)}|/|\alpha_2^{(1)}| = |\alpha_1^{(2)}|/|\alpha_2^{(2)}|$. On the other hand, the intensities of the dark parts of the two solitons s_1 and s_2 appearing in the third component remain unaltered after collision. Both bright and dark parts of the mixed solitons s_1 and s_2 experience a position-shift of same magnitude but of different signs. The position-shift experienced by soliton s_1 (and s_2) is $\Phi_1 = \frac{R_3-R_2-R_1}{2}$ (and $\Phi_2 = -\Phi_1$), where $R_1,\ R_2$ and R_3 are defined in Eq. (18) and the dark solitons s_1 and s_2 experience phase-shifts $2\phi_1^{(2)} - \pi$ and $-2\phi_1^{(1)} + \pi$, respectively. This shows that the bright components exhibit energy exchanging collisions characterized by an intensity redistribution (energy sharing) among the bright parts of the mixed solitons appearing in first two components and an amplitude dependent position-shift, whereas the dark parts of the two solitons undergo mere elastic collision accompanied by the same position shift as that of bright parts. Such a collision scenario is depicted in Fig. 5 for the parametric choice $k_1 = 2-i$, $k_2 = 1.5-i$, $\omega_1 = 1+i$, $\omega_2 = -1+i$, $m_1 = 0.7$, $\rho_1 = 4$, $b_1 = 2$, $\alpha_1^{(1)} = 1.2 + 0.2i$, $\alpha_2^{(1)} = -1 + 2i$, $\alpha_1^{(2)} = 0.25 + 0.25i$, and $\alpha_2^{(2)} = -1 + i$ at $t = 0$. In Fig. 5, the intensity of soliton s_1 is suppressed (enhanced) in the $S^{(1)}$ ($S^{(2)}$) component and the reverse occurs for soliton s_2. But the solitons appearing in $S^{(3)}$ and L components undergo elastic collision only.



FIG. 5: Energy exchanging collision of (2b-1d) mixed two-solitons in three-component LSRI system.

2. $(1b-2d)$ soliton collisions

Next we consider the collision scenario in the three-component LSRI system where the two colliding mixed solitons (say s_1 and s_2) are comprised of one bright part and two dark parts and appear respectively in the $S^{(1)}$ and $(S^{(2)}, S^{(3)})$ components. A careful asymptotic analysis of Eq. (18) with $m=1$ and $n=2$ shows that the amplitudes of the bright parts of the mixed solitons before and after collision are given by

$$\left(A_1^{1-},\ A_1^{2-},\ A_1^{1+},\ A_1^{2+}\right)=\left(\frac{\alpha_1^{(1)}}{2}e^{-\frac{R_1}{2}},\ \frac{1}{2}e^{\delta_{11}-(\frac{R_1+R_3}{2})},\ \frac{1}{2}e^{\delta_{21}-(\frac{R_2+R_3}{2})},\ \frac{\alpha_2^{(1)}}{2}e^{-\frac{R_2}{2}}\right).\quad(22)$$

Substitution of the expressions for various quantities from Eq. (18) with $m=1$ in Eq. (22) shows that the intensities of the bright parts of the two mixed solitons are same before and after interaction, i.e., $|A_1^{j+}|^2=|A_1^{j-}|^2,\ j=1,2$. Similarly, the amplitudes of the dark parts of the two mixed solitons before and after collision in the $S^{(2)}$ $(S^{(3)})$ component are same and are equal to ρ_l (ρ_2). This clearly indicates that the intensities of the dark parts of the colliding mixed solitons are unaltered during collision. Thus for the (1b-2d) case from the above expressions we observe that both the bright and dark parts of the mixed solitons undergo standard elastic collision of solitons accompanied by position-shifts of magnitude $|(R_3-R_2-R_1)/2|$, where R_1, R_2 and R_3 are defined in equation(18). The phase-shifts of the dark solitons s_1 and s_2 are $2\phi_1^{(2)}-\pi$ and $-2\phi_1^{(1)}+\pi$, respectively. Such an elastic collision behaviour is shown in Fig. 6 for $k_1=1-2i$, $k_2=1.5+i$, $\omega_1=-1-i$, $\omega_2=-2+i$, $m_1=-2$, $m_2=0.5$, $\rho_1=4$, $\rho_2=4$, $b_1=1$, $b_2=2$, $\alpha_1^{(1)}=0.5-i$, and $\alpha_2^{(1)}=1.4+i$ at $t=-3$. We do not present the plot for the long-wave component in Fig. 6 as it exhibits the collision process same as that of the $S^{(1)}$ component except for different amplitudes.



FIG. 6: Elastic collision of (1b-2d) mixed two-solitons in three-component LSRI system.

Our analysis reveals the interesting fact that the energy exchanging collision of mixed solitons can be realized only in the bright parts of the mixed solitons in three-component LSRI system and is possible only if the bright parts of the mixed solitons appear at least in two components.

The above analysis can be extended straightforwardly to arbitrary N-SW components case where the bright parts of the mixed soliton appear in m-components and the remaining $(N-m)$ components admit dark parts. It can be shown that the shape-changing (energy exchanging) collision is possible only if the bright parts of the mixed solitons appear at least in two short-wave components, i.e. $m \geq 2$.

VII. SOLITON BOUND STATES

Soliton bound states are another interesting class of multisoliton solutions. Soliton bound states can be viewed as composite solitons moving with a common speed. two-soliton bound states in the $(2+1)D$ two component LSRI system (12) with $q=2$ can be obtained from Eq. (18) for the choice $\frac{\omega_{1R}}{k_{1R}} = \frac{\omega_{2R}}{k_{2R}}$ and $k_{1I} = k_{2I}$, $m=1$, $n=1$, and the corresponding solution reads as

$$S^{(1)} = \frac{1}{D_1}\left(e^{\frac{l_1+\delta_{21}}{2}}\cosh(\hat{\eta}_{2R}+i\hat{\delta}_{21}) + e^{\frac{l_2+\delta_{11}}{2}}\cosh(\hat{\eta}_{1R}+i\hat{\delta}_{11})\right)e^{i\eta_{1I}}, \tag{23a}$$

$$\begin{aligned} S^{(2)} = \frac{\rho_1 e^{i\zeta_1}}{D_1}\Bigg[& e^{\frac{Q_3^{(1)}}{2}}\cosh\left(N_1 - i\frac{Q_{3I}^{(1)}}{2}\right) + e^{\frac{Q_{11}^{(1)}+Q_{22}^{(1)}}{2}}\cosh(N_2+i\hat{Q}_{11}) \\ & + e^{\frac{Q_{12}^{(1)}+Q_{21}^{(1)}}{2}}\cosh(N_3+i\hat{Q}_{12})\Bigg], \end{aligned} \tag{23b}$$

$$D_1 = e^{\frac{R_3}{2}}\cosh\left(\eta_{1R}+\eta_{2R}+\frac{R_3}{2}\right) + e^{\frac{R_1+R_2}{2}}\cosh\left(\eta_{1R}-\eta_{2R}+\frac{R_1-R_2}{2}\right) + e^{\frac{\delta_0+\delta_0^*}{2}}\cos(\delta_{0I}),$$

where $\eta_{1R} = k_{1R}\left[x + (2k_{1I} - \frac{\omega_{1R}}{k_{1R}})y + \frac{\omega_{1R}}{k_{1R}}t\right]$, $\eta_{2R} = \eta_{1R}(\frac{k_{2R}}{k_{1R}})$, $e^{l_1} = \alpha_1^{(1)}$, $e^{l_2} = \alpha_2^{(1)}$, $\hat{\delta}_{21} = \frac{\delta_{21I}-l_{1I}}{2}$, $\hat{\delta}_{11} = \frac{\delta_{11I}-l_{2I}}{2}$, $\hat{Q}_{11} = \frac{Q_{11I}^{(1)}-Q_{22I}^{(1)}}{2}$, $\hat{Q}_{12} = \frac{Q_{12I}^{(1)}-Q_{21I}^{(1)}}{2}$, $\hat{\eta}_{1R} = \eta_{1R} + \frac{\delta_{11R}-l_{2R}}{2}$, $\hat{\eta}_{2R} = \eta_{2R} + \frac{\delta_{21R}-l_{1R}}{2}$, $N_1 = \eta_{1R}+\eta_{2R}+\frac{Q_{3R}^{(1)}}{2}$, $N_2 = \eta_{1R}-\eta_{2R}+\frac{Q_{11R}^{(1)}-Q_{22R}^{(1)}}{2}$, $N_3 = \eta_{1I}-\eta_{2I}+\frac{Q_{12I}^{(1)}-Q_{21I}^{(1)}}{2}$, $\eta_{jI} = k_{jI}x - (k_{jR}^2 - k_{jI}^2 + \omega_{jI})y + \omega_{jI}t$, $j = 1, 2$, and all the other quantities found in the above expressions can be deduced from the corresponding quantities appearing in Eq. (18) by putting $m=1$ and $n=1$. The suffixes R and I appearing in the various quantities in Eq. (23) denote real and imaginary parts, respectively.

The two-soliton bound state is shown in Fig. 7 (top panel) for the choice $k_1 = 1+i$, $k_2 = 2+i$, $\omega_1 = -2-i$, $\omega_2 = -4+7i$, $m_1 = 0.7$, $t = -1$, $\alpha_1^{(1)} = 1+i$, $\alpha_2^{(1)} = 0.5+i$, $\rho_1 = 4$

and $b_1 = 2$. The bound state solitons display beating effects due to the oscillatory terms in Eq. (23). The beating effects can be suppressed completely by tuning the $\alpha_1^{(j)}$-parameters. This is shown in the bottom panel of Fig. 7, where the parameters are chosen same as that of the plots in the top panel except for $\alpha_1^{(1)}$, which is now fixed as 0.02. It should be noticed that though the α-parameters do not have any observable effects on the single soliton propagation in the two-SW components case as discussed in section IV A, they can display significant effects while considering bound state soliton propagation.



FIG. 7: Mixed two-soliton bound states in the two-component LSRI system. Top panel: with beating effects, Bottom panel: suppression of beating by tuning the $\alpha_1^{(j)}$-parameters.

Now it is of interest to investigate the influence of the ω_j, $j = 1, 2$, parameters on the bound state soliton dynamics which are arising due to the higher dimensional nature of the system. In this connection, we make $\omega_1 = \omega_2 = 0$ and we choose $k_{1I} = k_{2I}$. This will result in a bound state in the $x - y$ plane which is stationary in time. Such a two-soliton bound state with breathing oscillations in the $x - y$ plane is shown in the top panel of Fig. 8 for $\omega_1 = \omega_2 = 0$, $k_1 = 2 - i$, $k_2 = 1.5 - i$, $m_1 = 0.7$, $\rho_1 = 4$, $b_1 = 2$, $\alpha_1^{(1)} = 1.2 + 0.2i$, and $\alpha_2^{(1)} = 1 + 2i$ at $t = 0$. But when ω_1 and ω_2 become different and non-zero, they make the two solitons to undergo collision in the $x - y$ plane. Thus due to the presence of the ω_j-parameters and the higher dimensionality of the system there occurs a transition from bound states to interacting solitons and is shown in the bottom panel of Fig. 8. This also shows that the presence of ω_j parameters results in a wide range of parameters for which the soliton collision can take place.



FIG. 8: Mixed two-soliton bound state for $\omega_1 = \omega_2 = 0$ (top panel) and their transition to colliding solitons for $\omega_1 = 1 + i$, $\omega_2 = -1 + i$ (bottom panel) in two-component LSRI system. Other parameters are given in the text.

The dramatic change in the nature of soliton propagation in the $x - y$ plane due to the presence of ω_j-parameters resulting from the higher dimensional nature of the system (12) can exhibit additional features if we consider three SW components case. Soliton bound states in $x - y$ plane for $\omega_1 = \omega_2 = 0$, $k_1 = 2 - i$, $k_2 = 1.5 - i$, $m_1 = 0.7$, $\rho_1 = 4$, $b_1 = 2$, $\alpha_1^{(1)} = 1.2 + 0.2i$, $\alpha_2^{(1)} = -1 + 2i$, $\alpha_1^{(2)} = 0.25 + 0.25i$, and $\alpha_2^{(2)} = -1 + i$ at $t = 0$ is shown in Fig. 9. For the same parameters, with non-zero ω_j values, there occurs head-on collision of solitons and this is depicted in Fig. 5 for $\omega_1 = 1 + i$ and $\omega_2 = -1 + i$. Interestingly, the presence of second SW component now induces the fascinating collision involving energy exchange among the solitons in SW components. Also, the α-parameters can be tuned appropriately to suppress the beating effects of the bound soliton states.



FIG. 9: (2b-1d) mixed two-soliton bound states in three-component LSRI system for $\omega_1 = \omega_2 = 0$ and their transition to colliding solitons for $\omega_j \neq 0$ is given in Fig. 5.

VIII. CONCLUSION

In this paper, we have derived the multicomponent LSRI system for the propagation of weak nonlinear dispersive waves in $(2+1)D$. Then we consider the integrable multicomponent LSRI system in (2+1)-dimensions and obtain mixed (bright-dark) one- and two-soliton solutions using Hirota's method. Our study shows that the bright and dark parts of the mixed solitons in the 2-SW components behave like scalar bright and dark solitons. But in three-SW components case different types of splitting of the mixed soliton into bright and dark parts are possible which makes their dynamics interesting. Study on the collision dynamics of the mixed solitons shows that their collision in two-SW components case and in the three-SW components case with two dark parts and one bright part are elastic. However in the three-SW components case where the two colliding mixed solitons are split up into two bright parts and one dark part, the bright parts undergo energy exchanging collision characterized by intensity redistribution (energy sharing) and amplitude dependent position-shift. This collision process is also influenced by the soliton parameters of the dark part. The dark parts of the mixed solitons undergo only elastic collision.

Finally, we have considered the soliton bound states. To elucidate the understanding, we explicitly presented the two-soliton bound state expression for $m = 1$ and $n = 1$. Interestingly, we find that the α-parameters which do not show any significant effect on the one-soliton propagation display interesting effects on the bound states. Particularly, they can be profitably used in suppressing the beating effects. Another important observation follows from our above study is that the presence of ω_j-parameters can alter significantly the dynamics of solitons. Specifically, in the absence of ω_j-parameters, which can result due to the higher dimensional nature of the system, there occurs bound state solitons in the $x - y$ plane. But when the ω_j's are brought into picture the solitons exhibit collision behaviour. Physically, this means that due to the presence of ω_j-parameters the attractive force between the bound solitons vanishes and the solitons pass through each other. It has also been shown that for the three-SW components and one long-wave component case, for non-zero values of $\omega_j,\ j = 1, 2$, one can have energy exchanging collision for the bright parts of the mixed solitons in the $x - y$ plane whereas in the absence of ω_j's there occurs only stationary bound soliton in the $x - y$ plane.

Acknowledgement

T.K. acknowledges the support of the Department of Science and Technology, Government of India, in the form of a research project and also thank the principal and management of Bishop Heber College for constant support and encouragement. T.K. also thanks K. Sakkaravarthi for his assistance in the preparation of the manuscript. M.V. acknowledges the financial support from UGC-Dr. D. S. Kothari post-doctoral fellowship scheme. The work of M.L. is supported by a DST-IRPHA project. M.L. is also supported by a DST Ramanna Fellowship project and a DAE Ramanna Fellowship.

[1] Y. S. Kivshar and G. P. Agrawal, *Optical Solitons: From Fibers to Photonic Crystals* (Academic Press, San Diego, 2003).

[2] N. Akhmediev and A. Ankiewicz, *Solitons: Nonlinear Pulses and Beams* (Chapman and Hall, London, 1997).

[3] A. C. Scott, *Nonlinear Science: Emergence and Dynamics of Coherent Structures* (Oxford University Press, Oxford, 1999).

[4] M. Oikawa, M. Okamura, and M. Funakoshi, *J. Phys. Soc. Jpn.* **58**, 4416 (1989).

[5] R. H. J. Grimshaw, *Stud. Appl. Math.* **56**, 241 (1977).

[6] M. Boiti, J. Lenon, F. Pempinelli, and A. S. Pire, *J. Phys. A: Math. Gen.* **37**, 4243 (2004).

[7] V. E. Zakharov, *Zh. Eksp. Teor. Fiz.* **62**, 1745 (1972) [*Sov. Phys. JETP* **35**, 908 (1972)]

[8] Y. S. Kivshar, *Opt. Lett.* **17**, 1322 (1992).

[9] N. Yajima and M. Oikawa, *Prog. Theor. Phys.* **56**, 1719 (1976).

[10] Y. C. Ma, *Stud. Appl. Math.* **59**, 201 (1978).

[11] Y. C. Ma and L. G. Redekopp, *Phys. Fluids* **22**, 1872 (1979).

[12] M. Funakoshi and M. Oikawa, *J. Phys. Soc. Jpn.* **52**, 1982 (1983).

[13] D. J. Benney, *Stud. Appl. Math.* **55**, 93 (1976).

[14] Y. Ohta, K. Maruno, and M. Oikawa, *J. Phys. A: Math. Theor.* **40**, 7659 (2007).

[15] H. E. Nistazakis, D. J. Frantzeskakis, P. G. Kevrekidis, B. A. Malomed, and R. Carretero-Gonzalez, *Phys. Rev. A* **77**, 033612 (2008).

[16] A. Visinescu, D. Grecu, R. Fedele, and S. De Nicola, *SIGMA* **7**, 041 (2011).

[17] T. Kanna, M. Vijayajayanthi, K. Sakkaravarthi, and M. Lakshmanan, *J. Phys. A: Math. Theor.* **42**, 115103 (2009).

[18] M. H. Jakubowski, K. Steiglitz, and R. Squier, *Phys. Rev. E* **58**, 6752 (1998).

[19] K. Steiglitz, *Phys. Rev. E* **63**, 016608 (2000).

[20] T. Kanna and M. Lakshmanan, *Phys. Rev. E* **67**, (2003) 046617.

[21] A. P. Sheppard and Y. S. Kivshar, *Phys. Rev. E* **55**, 4773 (1997).

[22] Y. Ohta, D. Wang, and J. Yang, *Stud. Appl. Math.* **127**, 345 (2011).

[23] M. Vijayajayanthi, T. Kanna, and M. Lakshmanan, *Phys. Rev. A* **77**, 013820 (2008).

[24] R. Hirota, *The Direct Method in Soliton Theory* (Cambridge University Press, Cambridge, 2004).

[25] J. Hietarinta, *J. Math. Phys.* **28**, 1732 (1987), *ibid* **28**, 2094 (1987); *ibid* **28**, 2586 (1987); J. Hietarinta, *J. Math. Phys.* **29**, 628 (1988).

[26] T. Yamagata, *J. Meteor. Soc. Jpn.* **58**, 160 (1980).

[27] C. G. Koop and L. G. Redekopp, *J. Fluid Mech.* **111**, 367 (1981).